



KIIT POLYTECHNIC

LECTURE NOTES

ON

ENGG. MATH -I

PART-3

Prepared by

Satyajit Mohapatra

(Lecturer)

Department of Basic Sciences and Humanities, KIIT Polytechnic BBSR

CONTENTS

S.No	Chapter Name	Page No
1	Trigonometry	3-20

CHAPTER-5

TRIGONOMETRY

Trigonometry is a branch of mathematics that studies triangles and the relationships between the lengths of their sides and the angles.


It is derived from three small Greek words **tri + gono+ metry**

Tri---- three,

Gono---- angle,

Metry----- measurement

Angle:An angle can be formed by the union of two rays or two lines with a common point(called vertex).



1. If an angle measured in anticlockwise direction then it is positive .
2. If an angle measured in clockwise direction then it is negative.

Angles are measured using the following units:

1. Sexagesimal degree ($^{\circ}$)

If the circumference of a circle is divided into 360 equal parts, the central angle corresponding to each of its 360 parts is an angle of one degree sexagesimal, (1°).

A **degree** has **60 minutes ('')** [$1^{\circ} = 60'$]and a **minute** has **60 seconds (")** [$1' = 60''$].


2. Radian (rad):It is the measure of an angle whose arc is a ratio.

$$2\pi \text{ radian} = 360^{\circ} \quad \text{and } \pi \text{ radian} = 180^{\circ}$$

$$1 \text{ radian} = \frac{180^{\circ}}{\pi} \quad \text{and } 1^{\circ} = \frac{\pi}{180} \text{ radian}$$

Figure

Radian measurement



Conversion from Degree to Radian:

$$30^\circ = 30 \times \frac{\pi}{180} = \frac{\pi}{6}$$

$$60^\circ = 60 \times \frac{\pi}{180} = \frac{\pi}{3}$$

$$90^\circ = 90 \times \frac{\pi}{180} = \frac{\pi}{2}$$

$$150^\circ = 150 \times \frac{\pi}{180} = \frac{5\pi}{6}$$

$$270^\circ = 270 \times \frac{\pi}{180} = \frac{3\pi}{2}$$

Conversion from Radian to Degree:


$$\frac{\pi}{2} = \frac{\pi}{2} \times \frac{180}{\pi} = 90^\circ$$

Trigonometric Ratios:

It is the ratio of the sides of a right angled triangle with respect to base angle.

We have six trigonometric ratios like sine, cosine, tangent, cotangent, secant and cosecant.(i.e sin, cos, tan ,cot ,sec and cosec).

How to find all trigonometric ratios from a right angled triangle.



In a right triangle, the following trigonometric ratios can be defined:

Let $\angle B = \theta$, $\angle A = 90^\circ$, and $\angle C = 90 - \theta$,

Let AB = c = Base, AC = b = Perpendicular and BC = a = Hypotenuse.

Sine

The **sine** of the angle θ is the ratio between the length of the **opposite side(Perpendicular)** and the **hypotenuse** of the triangle.

It is denoted by **sin θ**

$$\sin\theta = \frac{\text{perpendicular}}{\text{hypotenuse}} = \frac{b}{a} \text{ or } \frac{p}{h}$$

Cosine

The **cosine** of the angle θ is the **ratio** between the length of the **adjacent side(base)** and the **hypotenuse** of the triangle.

It is denoted by **cos θ** .

$$\cos\theta = \frac{\text{base}}{\text{hypotenuse}} = \frac{c}{a} \text{ or } \frac{b}{h}$$

Tangent

The **tangent** of the angle θ is the **ratio** between the length of the **opposite side(perpendicular)** and the **adjacent side(base)** of the triangle.

It is denoted by **tan θ** .

$$\tan\theta = \frac{b}{c} \text{ or } \frac{p}{b}$$

Cotangent

The **cotangent** of the angle θ is the **inverse** of the **tangent** of θ .

It is denoted by **cot θ** .

$$\cot\theta = \frac{c}{b} \text{ or } \frac{b}{p}$$

Secant

The **secant** of angle θ is the **inverse** of the **cosine** of θ .

It is denoted by **sec θ** .

$$\sec\theta = \frac{a}{c} \text{ or } \frac{h}{b}$$

Cosecant

The **cosecant** of angle θ is the **inverse** of the **sine** of θ .

It is denoted by **cosec θ** .

$$\cosec\theta = \frac{a}{b} \text{ or } \frac{h}{p}$$

Formula 1:

$$\sin\theta \cosec\theta = 1, \sin\theta = \frac{1}{\cosec\theta}, \cosec\theta = \frac{1}{\sin\theta}$$

$$\tan\theta \cot\theta = 1, \tan\theta = \frac{1}{\cot\theta}, \cot\theta = \frac{1}{\tan\theta}$$

$$\cos\theta \sec\theta = 1, \cos\theta = \frac{1}{\sec\theta}, \sec\theta = \frac{1}{\cos\theta}$$

Formula 2:

$$\sin^2\theta + \cos^2\theta = 1, \sin^2\theta = 1 - \cos^2\theta, \cos^2\theta = 1 - \sin^2\theta$$

$$1 + \tan^2\theta = \sec^2\theta, \sec^2\theta - \tan^2\theta = 1, \sec^2\theta - 1 = \tan^2\theta$$

$$1 + \cot^2\theta = \cosec^2\theta, \cosec^2\theta - \cot^2\theta = 1, \cosec^2\theta - 1 = \cot^2\theta$$

Formula 3: $\tan\theta = \frac{\sin\theta}{\cos\theta}$, $\cot\theta = \frac{\cos\theta}{\sin\theta}$

Value of the trigonometric ratios in different angles:

a:	0°	30°	45°	60°	90°	180°	270°
\sin	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1	0	-1
\cos	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0	-1	0
\tan	0	$\frac{\sqrt{3}}{3}$	1	$\sqrt{3}$	$\rightarrow \infty$	0	$\rightarrow -\infty$

Define all trigonometric ratios at different angles:

$\sin(90^\circ - \theta) = \cos\theta$
 $\cos(90^\circ - \theta) = \sin\theta$
 $\tan(90^\circ - \theta) = \cot\theta$
 $\cot(90^\circ - \theta) = \tan\theta$
 $\sec(90^\circ - \theta) = \cosec\theta$
 $\cosec(90^\circ - \theta) = \sec\theta$

$\sin(180^\circ - \theta) = \sin\theta$
 $\cos(180^\circ - \theta) = -\cos\theta$
 $\tan(180^\circ - \theta) = -\tan\theta$
 $\cot(180^\circ - \theta) = -\cot\theta$
 $\sec(180^\circ - \theta) = -\sec\theta$
 $\cosec(180^\circ - \theta) = \cosec\theta$

$\sin(180^\circ + \theta) = -\sin\theta$
 $\cos(180^\circ + \theta) = -\cos\theta$
 $\tan(180^\circ + \theta) = \tan\theta$
 $\cot(180^\circ + \theta) = \cot\theta$
 $\sec(180^\circ + \theta) = -\sec\theta$
 $\cosec(180^\circ + \theta) = -\cosec\theta$

$\sin(-\theta) = -\sin\theta$
 $\cos(-\theta) = \cos\theta$
 $\tan(-\theta) = -\tan\theta$
 $\cot(-\theta) = -\cot\theta$
 $\sec(-\theta) = \sec\theta$
 $\cosec(-\theta) = -\cosec\theta$

$$\begin{aligned}\sin(90^\circ + \theta) &= \cos\theta \\ \cos(90^\circ + \theta) &= -\sin\theta \\ \tan(90^\circ + \theta) &= -\cot\theta \\ \cot(90^\circ + \theta) &= -\tan\theta \\ \sec(90^\circ + \theta) &= -\cosec\theta \\ \cosec(90^\circ + \theta) &= \sec\theta\end{aligned}$$

$$\begin{aligned}\sin(270^\circ + \theta) &= -\cos\theta \\ \cos(270^\circ + \theta) &= \sin\theta \\ \tan(270^\circ + \theta) &= -\cot\theta \\ \cot(270^\circ + \theta) &= -\tan\theta \\ \sec(270^\circ + \theta) &= \cosec\theta \\ \cosec(270^\circ + \theta) &= -\sec\theta\end{aligned}$$

$$\begin{aligned}\sin(270^\circ - \theta) &= -\cos\theta \\ \cos(270^\circ - \theta) &= -\sin\theta \\ \tan(270^\circ - \theta) &= \cot\theta \\ \cot(270^\circ - \theta) &= \tan\theta \\ \sec(270^\circ - \theta) &= -\cosec\theta \\ \cosec(270^\circ - \theta) &= -\sec\theta\end{aligned}$$

$$\begin{aligned}\sin(360^\circ - \theta) &= -\sin\theta \\ \cos(360^\circ - \theta) &= \cos\theta \\ \tan(360^\circ - \theta) &= -\tan\theta \\ \cot(360^\circ - \theta) &= -\cot\theta \\ \sec(360^\circ - \theta) &= \sec\theta \\ \cosec(360^\circ - \theta) &= -\cosec\theta\end{aligned}$$

$$\begin{aligned}\sin(360^\circ + \theta) &= \sin\theta \\ \cos(360^\circ + \theta) &= \cos\theta \\ \tan(360^\circ + \theta) &= \tan\theta \\ \cot(360^\circ + \theta) &= \cot\theta \\ \sec(360^\circ + \theta) &= \sec\theta \\ \cosec(360^\circ + \theta) &= \cosec\theta\end{aligned}$$


$$\sin(n\pi + \theta) = (-1)^n \sin \theta, \sin(n\pi + \theta) = \begin{cases} \sin \theta, & \text{if } n \text{ is even} \\ -\sin \theta, & \text{if } n \text{ is odd} \end{cases}$$

$$\cos(n\pi + \theta) = (-1)^n \cos \theta, \cos(n\pi + \theta) = \begin{cases} \cos \theta, & \text{if } n \text{ is even} \\ -\cos \theta, & \text{if } n \text{ is odd} \end{cases}$$

$\tan(n\pi + \theta) = \tan \theta$, if n is either even or odd. Where n is an integer.

Similarly we can find cot, sec and cosec

Let $0 < \theta < 90^\circ$



A \Rightarrow In the 1st quadrant all trigonometric ratios are positive

S \Rightarrow In the 2nd quadrant only sine and cosec are positive but rest are negative

T \Rightarrow In the 3rd quadrant only tan and cot are positive but rest are negative

C \Rightarrow In the 4th quadrant only cos and sec are positive but rest are negative

Ex: Show which of the following are positive.

$$\sin 271^\circ, \cos 375^\circ, \tan 99^\circ, \cot 333^\circ, \sec 159^\circ \text{ and } \cosec 79^\circ$$

Ex: Express the following trigonometric ratios in the form of acute angle.

$$\sin 271^\circ, \cos 375^\circ, \tan 333^\circ, \cot 1233^\circ, \sec (-2349)^\circ \text{ and } \cosec 1230^\circ$$

Ex: find the value of $\sin 1230^\circ, \cos 210^\circ, \tan 900^\circ \text{ and } \cot 750^\circ$

Ex: Find the value of A if $\cot 3A = \tan 6A$

Ex: Find the value of $\cos 1 \cdot \cos 2 \cdot \cos 3 \dots \cos 100$.

Ex: Find the value of $\sin 1 \cdot \sin 2 \cdot \sin 3 \dots \sin 200$.

Ex: Find the value $\tan 1 \cdot \tan 2 \cdot \tan 3 \dots \tan 89$.

Ex: find the value of $\tan 9 \cdot \tan 27 \cdot \tan 45 \cdot \tan 63 \cdot \tan 81$.

Ex: Find the value of $\log \sin 1 \cdot \log \sin 2 \cdot \log \sin 3 \dots \log \sin 100$.

Ex: Find the value of $\log \tan 1 + \log \tan 2 + \dots + \log \tan 89$.

Ex: If $\sin A = \frac{3}{5}$ and $90^\circ < A < 180^\circ$ then find other trigonometric ratios.

Ans: Given $\sin A = \frac{3}{5}$

$$\cos A = \sqrt{1 - \sin^2 A} = \sqrt{1 - \left(\frac{3}{5}\right)^2} = \pm \frac{4}{5}$$

Due to $90^\circ < A < 180^\circ$ $\cos A$ is negative. So $\cos A = -\frac{4}{5}$, similarly

$$\tan A = \frac{\sin A}{\cos A} = -\frac{3}{4}, \cot A = \frac{1}{\tan A} = -\frac{4}{3}, \sec A = \frac{1}{\cos A} = -\frac{5}{4}, \cosec A = \frac{1}{\sin A} = \frac{5}{3}$$

Ex: If $\cos A = \frac{4}{5}$ and $180^\circ < A < 270^\circ$ then find the value of other trigonometric ratios

Trigonometric ratios of compound angles, multiple angles and sub-multiple angles

Compound Angles: The sum or difference of two or more angles is known as multiple angles.

Ex: $A+B, A-B, A+B+C \dots$

Multiple Angles: If an angle A is multiplied by a non zero integer then it is called multiple angles.

Ex: $2A, 3A, \dots$

Sub-multiple Angles: If an angle A is divided by a non zero integer then it is called sub-multiple angles.

Ex: $\frac{A}{2}, \frac{A}{3}, \dots$

Formulas:

Formula-1

$$\sin(A + B) = \sin A \cos B + \cos A \sin B$$

$$\sin(A - B) = \sin A \cos B - \cos A \sin B$$

$$\cos(A + B) = \cos A \cos B - \sin A \sin B$$

$$\cos(A - B) = \cos A \cos B + \sin A \sin B$$

Formula-3

$$\sin(A + B) + \sin(A - B) = 2 \sin A \cos B$$

$$\sin(A + B) - \sin(A - B) = 2 \cos A \sin B$$

$$\cos(A + B) + \cos(A - B) = 2 \cos A \cos B$$

$$\{\cos(A + B) - \cos(A - B) = -2 \sin A \sin B$$

$$\text{or } \cos(A - B) - \cos(A + B) = 2 \sin A \sin B$$

Formula-5

$$\sin(A + B) \sin(A - B) = \begin{cases} \sin^2 A - \sin^2 B \\ \cos^2 B - \cos^2 A \end{cases}$$

$$\cos(A + B) \cos(A - B) = \begin{cases} \cos^2 A - \sin^2 B \\ \cos^2 B - \sin^2 A \end{cases}$$

Formula-2

$$\tan(A + B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}$$

$$\tan(A - B) = \frac{\tan A - \tan B}{1 + \tan A \tan B}$$

$$\cot(A \pm B) = \frac{\cot A \cot B \mp 1}{\cot B \pm \cot A}$$

Formula-4

$$\sin C + \sin D = 2 \sin \frac{C+D}{2} \cos \frac{C-D}{2}$$

$$\sin C - \sin D = 2 \cos \frac{C+D}{2} \sin \frac{C-D}{2}$$

$$\cos C + \cos D = 2 \cos \frac{C+D}{2} \cos \frac{C-D}{2}$$

$$\cos C - \cos D = \begin{cases} -2 \sin \frac{C+D}{2} \sin \frac{C-D}{2} \\ 2 \sin \frac{C+D}{2} \sin \frac{D-C}{2} \end{cases}$$

Ex: Find the value of $\sin 15^\circ$, $\cos 15^\circ$, $\tan 15^\circ$ and $\cot 15^\circ$.

$$\text{Ans: } \sin 15^\circ = \sin(45^\circ - 30^\circ) = \sin 45^\circ \cos 30^\circ - \cos 45^\circ \sin 30^\circ$$

$$= \frac{1}{\sqrt{2}} \times \frac{\sqrt{3}}{2} - \frac{1}{\sqrt{2}} \times \frac{1}{2} = \frac{\sqrt{3}}{2\sqrt{2}} - \frac{1}{2\sqrt{2}} = \frac{\sqrt{3}-1}{2\sqrt{2}}$$

$$\cos 15^\circ = \cos(45^\circ - 30^\circ) = \cos 45^\circ \cos 30^\circ + \sin 45^\circ \sin 30^\circ$$

$$= \frac{1}{\sqrt{2}} \times \frac{\sqrt{3}}{2} + \frac{1}{\sqrt{2}} \times \frac{1}{2} = \frac{\sqrt{3}}{2\sqrt{2}} + \frac{1}{2\sqrt{2}} = \frac{\sqrt{3}+1}{2\sqrt{2}}$$

$$\tan 15^\circ = \tan(45^\circ - 30^\circ) = \frac{\tan 45^\circ - \tan 30^\circ}{1 + \tan 45^\circ \tan 30^\circ} = \frac{1 - \frac{1}{\sqrt{3}}}{1 + 1 \times \frac{1}{\sqrt{3}}} = \frac{\sqrt{3}-1}{\sqrt{3}+1}$$

Or we can derive $\tan 15^\circ = \frac{\sin 15^\circ}{\cos 15^\circ}$

$$\cot 15^\circ = \frac{\sqrt{3}+1}{\sqrt{3}-1}$$

Ex: Find the value of $\sin 75^\circ$, $\cos 75^\circ$, $\tan 75^\circ$ and $\cot 75^\circ$

Hints: Expand $75^\circ = 45^\circ + 30^\circ$, use the formula you will get the answer.

Ex: if $\tan A = \frac{1}{2}$ and $\tan B = \frac{1}{3}$ then find the value of $A + B$.

Ans: Given $\tan A = \frac{1}{2}$ and $\tan B = \frac{1}{3}$, use $\tan(A + B)$ formula

$$\tan(A + B) = \frac{\frac{1}{2} + \frac{1}{3}}{1 - \frac{1}{2} \times \frac{1}{3}} = \frac{\frac{5}{6}}{\frac{5}{6}} = 1 = \tan 45^\circ \Rightarrow A + B = 45^\circ$$

Ex: If $\tan A = \frac{13}{27}$ and $\tan B = \frac{7}{20}$ then find $A + B$.

Ex: Show that $\sin 10^\circ + \sin 50^\circ - \sin 70^\circ = 0$

$$\begin{aligned} \text{Ans: L.H.S } \sin 10^\circ + \sin 50^\circ - \sin 70^\circ &= \sin 10^\circ + \sin(60^\circ - 10^\circ) - \sin(60^\circ + 10^\circ) \\ &= \sin 10^\circ - 2\cos 60^\circ \sin 10^\circ = \sin 10^\circ - \sin 10^\circ = 0 \end{aligned}$$

Formula-6

$$\begin{aligned} \sin 2A &= 2\sin A \cos A \\ \cos 2A &= \begin{cases} \cos^2 A - \sin^2 A \\ 2\cos^2 A - 1 \\ 1 - 2\sin^2 A \end{cases} \end{aligned}$$

Formula-7

$$\begin{aligned} \sin 2A &= \frac{2\tan A}{1 + \tan^2 A} \\ \cos 2A &= \frac{1 - \tan^2 A}{1 + \tan^2 A} \\ \tan 2A &= \frac{2\tan A}{1 - \tan^2 A} \end{aligned}$$

Formula-8

$$\begin{aligned} \sin 3A &= 3\sin A - 4\sin^3 A \\ \cos 3A &= 4\cos^3 A - 3\cos A \\ \tan 3A &= \frac{3\tan A - \tan^3 A}{1 - 3\tan^2 A} \end{aligned}$$

Formula-9

$$\begin{aligned} \sin^2 A &= \frac{1 - \cos 2A}{2} \\ \cos^2 A &= \frac{1 + \cos 2A}{2} \\ \tan^2 A &= \frac{1 - \cos 2A}{1 + \cos 2A} \end{aligned}$$

Formula-10

$$\begin{aligned} \sin A &= 2\sin \frac{A}{2} \cos \frac{A}{2} \text{ or } \frac{2\tan \frac{A}{2}}{1 + \tan^2 \frac{A}{2}} \\ \cos A &= \cos^2 \frac{A}{2} - \sin^2 \frac{A}{2} \text{ or } 2\cos^2 \frac{A}{2} - 1 \text{ or } 1 - 2\sin^2 \frac{A}{2} \text{ or } \frac{1 - \tan^2 \frac{A}{2}}{1 + \tan^2 \frac{A}{2}} \end{aligned}$$

Formula-11

$$1 + \sin 2A = (\cos A + \sin A)^2$$

$$1 - \sin 2A = (\cos A - \sin A)^2$$

$$1 + \sin A = \left(\cos \frac{A}{2} + \sin \frac{A}{2} \right)^2$$

$$1 - \sin A = \left(\cos \frac{A}{2} - \sin \frac{A}{2} \right)^2$$

Ex: find the value of $\sin 105^\circ \cdot \cos 105^\circ$

$$\text{Ans: } \sin 105^\circ \cdot \cos 105^\circ = \frac{2\sin 105^\circ \cdot \cos 105^\circ}{2} = \frac{\sin 210^\circ}{2} = \frac{\sin(180^\circ + 30^\circ)}{2} = \frac{-\sin 30^\circ}{2} = -\frac{1}{4}$$

Ex: Show that $\tan 75^\circ + \cot 75^\circ = 4$

$$\text{Ans: L.H.S } \tan 75^\circ + \cot 75^\circ = \frac{\sin 75^\circ}{\cos 75^\circ} + \frac{\cos 75^\circ}{\sin 75^\circ} = \frac{\sin^2 75^\circ + \cos^2 75^\circ}{\sin 75^\circ \cos 75^\circ} = \frac{1}{\sin 75^\circ \cos 75^\circ} = \frac{2}{2\sin 75^\circ \cos 75^\circ} = \frac{2}{\sin 150^\circ} = \frac{2}{\left(\frac{1}{2}\right)} = 4$$

Note : The value of $\sin 18^\circ = \frac{\sqrt{5}-1}{4}$ and $\cos 36^\circ = \frac{\sqrt{5}+1}{4}$

Ex: Find the value of $\sin^2 18^\circ + \cos^2 36^\circ$.

$$\text{Ans: } \sin^2 18^\circ + \cos^2 36^\circ = \left(\frac{\sqrt{5}-1}{4}\right)^2 + \left(\frac{\sqrt{5}+1}{4}\right)^2 = \frac{(\sqrt{5}-1)^2 + (\sqrt{5}+1)^2}{16} = \frac{2(5+1)}{16} = \frac{3}{4}$$

Ex: Prove that $\frac{\cos 9^\circ + \sin 9^\circ}{\cos 9^\circ - \sin 9^\circ} = \tan 54^\circ$

$$\text{Ans: L.H.S } \frac{\cos 9^\circ + \sin 9^\circ}{\cos 9^\circ - \sin 9^\circ} = \frac{\frac{\cos 9^\circ}{\cos 9^\circ} + \frac{\sin 9^\circ}{\cos 9^\circ}}{\frac{\cos 9^\circ}{\cos 9^\circ} - \frac{\sin 9^\circ}{\cos 9^\circ}} = \frac{1 + \tan 9^\circ}{1 - \tan 9^\circ} = \frac{\tan 45^\circ + \tan 9^\circ}{1 - \tan 45^\circ \tan 9^\circ} = \tan(45^\circ + 9^\circ) = \tan 54^\circ$$

Ex: Prove that $\sec 2A + \tan 2A = \tan\left(\frac{\pi}{4} + A\right)$

$$\text{Ans: L.H.S } \sec 2A + \tan 2A = \frac{1}{\cos 2A} + \frac{\sin 2A}{\cos 2A} = \frac{1 + \sin 2A}{\cos 2A} = \frac{(\cos A + \sin A)^2}{\cos^2 A - \sin^2 A} = \frac{\cos A + \sin A}{\cos A - \sin A}$$

divide $\cos A$ in both N^r and D^r

$$= \frac{1 + \tan A}{1 - \tan A} = \frac{\tan \frac{\pi}{4} + \tan A}{1 - \tan \frac{\pi}{4} \tan A} = \tan\left(\frac{\pi}{4} + A\right)$$

Formula-12

$$\tan \frac{A}{2} = \frac{1 - \cos A}{\sin A} \text{ or } \frac{\sin A}{1 + \cos A}$$

$$\cot \frac{A}{2} = \frac{1 + \cos A}{\sin A} \text{ or } \frac{\sin A}{1 - \cos A}$$

Ex: find the value of $\sin 105^\circ \cdot \cos 105^\circ$

$$\text{Ans: } \sin 105^\circ \cdot \cos 105^\circ = \frac{2\sin 105^\circ \cdot \cos 105^\circ}{2} = \frac{\sin 210^\circ}{2} = \frac{\sin(180^\circ + 30^\circ)}{2} = \frac{-\sin 30^\circ}{2} = -\frac{1}{4}$$

Ex: Show that $\tan 75^\circ + \cot 75^\circ = 4$

$$\text{Ans: L.H.S } \tan 75^\circ + \cot 75^\circ = \frac{\sin 75^\circ}{\cos 75^\circ} + \frac{\cos 75^\circ}{\sin 75^\circ} = \frac{\sin^2 75^\circ + \cos^2 75^\circ}{\sin 75^\circ \cos 75^\circ} = \frac{1}{\sin 75^\circ \cos 75^\circ} = \frac{2}{2\sin 75^\circ \cos 75^\circ} = \frac{2}{\sin 150^\circ} = \frac{2}{\left(\frac{1}{2}\right)} = 4$$

Note : The value of $\sin 18^\circ = \frac{\sqrt{5}-1}{4}$ and $\cos 36^\circ = \frac{\sqrt{5}+1}{4}$

Ex: Find the value of $\sin^2 18^\circ + \cos^2 36^\circ$.

$$\text{Ans: } \sin^2 18^\circ + \cos^2 36^\circ = \left(\frac{\sqrt{5}-1}{4}\right)^2 + \left(\frac{\sqrt{5}+1}{4}\right)^2 = \frac{(\sqrt{5}-1)^2 + (\sqrt{5}+1)^2}{16} = \frac{2(5+1)}{16} = \frac{3}{4}$$

Ex: Prove that $\frac{\cos 9^\circ + \sin 9^\circ}{\cos 9^\circ - \sin 9^\circ} = \tan 54^\circ$

$$\text{Ans: L.H.S } \frac{\cos 9^\circ + \sin 9^\circ}{\cos 9^\circ - \sin 9^\circ} = \frac{\frac{\cos 9^\circ}{\cos 9^\circ} + \frac{\sin 9^\circ}{\cos 9^\circ}}{\frac{\cos 9^\circ}{\cos 9^\circ} - \frac{\sin 9^\circ}{\cos 9^\circ}} = \frac{1 + \tan 9^\circ}{1 - \tan 9^\circ} = \frac{\tan 45^\circ + \tan 9^\circ}{1 - \tan 45^\circ \tan 9^\circ} = \tan(45^\circ + 9^\circ) = \tan 54^\circ$$

Ex: Prove that $\sec 2A + \tan 2A = \tan\left(\frac{\pi}{4} + A\right)$

$$\text{Ans: L.H.S } \sec 2A + \tan 2A = \frac{1}{\cos 2A} + \frac{\sin 2A}{\cos 2A} = \frac{1 + \sin 2A}{\cos 2A} = \frac{(\cos A + \sin A)^2}{\cos^2 A - \sin^2 A} = \frac{\cos A + \sin A}{\cos A - \sin A}$$

divide $\cos A$ in both N^r and D^r

$$= \frac{1 + \tan A}{1 - \tan A} = \frac{\tan \frac{\pi}{4} + \tan A}{1 - \tan \frac{\pi}{4} \tan A} = \tan\left(\frac{\pi}{4} + A\right)$$

Assignment-1

Prove that

$$1. \frac{\tan\theta}{1+\tan^2\theta} = \sin\theta \cos\theta$$

$$2. \frac{1-\tan^2\theta}{1+\tan^2\theta} = \cos^2\theta - \sin^2\theta$$

$$3. \tan\theta + \cot\theta = \cosec\theta \sec\theta$$

$$4. (1 - \cos^2\theta) \cosec^2\theta = 1$$

$$5. \sec\theta \sqrt{(1 - \sin^2\theta)} = 1$$

$$6. \tan\theta \sin\theta + \cos\theta = \sec\theta$$

$$7. (1 - \cos\theta)(1 + \cos\theta)(1 + \cot^2\theta) = 1$$

$$8. \frac{\cos\theta}{1-\tan\theta} + \frac{\sin\theta}{1-\cot\theta} = \sin\theta + \cos\theta$$

$$9. (1 - \sin\theta)(1 + \cosec\theta) = \cos\theta \cot\theta$$

$$10. \sin\theta \cos\theta \tan\theta = \sin^2\theta$$

$$11. \sec^4\theta - \tan^4\theta = 2\sec^2\theta - 1$$

$$12. \sin\theta + \sin\theta \cot^2\theta = \cosec\theta$$

$$13. \frac{1}{\cosec A - \cot A} + \frac{1}{\cosec A + \cot A} = \frac{2}{\sin A}$$

$$14. \frac{\tan\theta + \sec\theta - 1}{\tan\theta - \sec\theta + 1} = \frac{1 + \sin\theta}{\cos\theta}$$

Assignment-2

$$15. \frac{\cot\theta + \cosec\theta - 1}{\cot\theta - \cosec\theta + 1} = \frac{1 + \cos\theta}{\sin\theta}$$

$$16. \sqrt{\frac{1+\sin\theta}{1-\sin\theta}} = \sec\theta + \tan\theta$$

$$17. \sqrt{\frac{1-\cos\theta}{1+\cos\theta}} = \cosec\theta - \cot\theta$$

$$18. \frac{\sin\theta - \cos\theta}{\sin\theta - \cos\theta} = \frac{\cos 2\theta}{1 + \sin 2\theta}$$

$$19. \frac{\sin\theta}{1-\cos\theta} + \frac{1-\cos\theta}{\sin\theta} = 2\cosec\theta$$

$$20. \sec^4 A - \sec^2 A = \tan^2 A + \tan^4 A$$

$$21. (1 + \cot\theta - \cosec\theta)(1 + \tan\theta + \sec\theta) = 2$$

$$22. (\cosec\theta - \sin\theta)(\sec\theta - \cos\theta)(\tan\theta + \cot\theta) = 1$$

$$23. \frac{\sin\theta \sec\theta}{\cos^2\theta} = \tan^3\theta + \tan\theta$$

$$24. \frac{1-\sin A}{1+\sin A} = (\sec A - \tan A)^2$$

$$25. \sqrt{\frac{\sec A - 1}{\sec A + 1}} = \cosec A - \cot A$$

Assignment-3

Prove that

$$1. \frac{1+\cos\theta+\sin\theta}{1+\cos\theta-\sin\theta} = \frac{\cos\theta}{1-\sin\theta}$$

$$2. \sin^6 A + \cos^6 A = 1 - 3\sin^2 A \cos^2 A$$

$$3. \frac{\tan A}{1-\cot A} + \frac{\cot A}{1-\tan A} = \sec A \cosec A + 1$$

$$4. (\sec\theta - 1) \cosec\theta - (1 - \cos\theta) \cot\theta = \tan\theta - \sin\theta$$

5. If $\cos\theta = -\frac{1}{2}$ and $180^\circ < \theta < 270^\circ$ then find all the trigonometric ratios.

$$6. \frac{\cos(180^\circ - A) \cot(90^\circ + A) \cos(-A)}{\tan(180^\circ + A) \tan(270^\circ + A) \sin(360^\circ - A)} = \cos A$$

$$7. \frac{\cos(180^\circ - A) \sin(90^\circ + A) \tan(90^\circ + A)}{\tan(270^\circ + A) \sin(270^\circ - A) \sin(90^\circ - A)} = 1$$

$$8. \frac{\cot(180^\circ + A) \cot(90^\circ + A) \cos(360^\circ - A)}{\tan(90^\circ + A) \cosec(180^\circ - A) \sin(-A)} = \sin A$$

Assignment:4

1. Find the value of $\cos 210^\circ$, $\sin(-1125^\circ)$, $\tan 480^\circ$

2. Express as the trigonometric ratio of an acute angle

$\tan 1129^\circ$, $\cot 1239^\circ$, $\cosec 3333^\circ$, $\cos(-439^\circ)$, $\sin(-1339^\circ)$

3. Prove that $\tan 225^\circ \cot 405^\circ + \tan 765^\circ \cot 675^\circ = 0$

4. Prove that $\tan 720^\circ - \cot 270^\circ - \sin 150^\circ \cos 120^\circ = \frac{1}{4}$

5. Prove that $\cos^2 \frac{\pi}{6} + \cosec \frac{5\pi}{6} + 3\tan^2 \frac{\pi}{6} = 6$

6. Prove that $\frac{2\sin\theta \cos\theta - \sin\theta}{1 - \sin\theta + \sin^2\theta - \cos^2\theta} = \cot\theta$

7. $\cos\theta(\tan\theta + 2)(2\tan\theta + 1) = 2\sec\theta + 5\sin\theta$

Assignment-5

Ex: Find the value of $\tan 105^\circ$ and $\tan \frac{13\pi}{12}$

Ex: Prove that $\sin(n+1)A\sin(n+2)A + \cos(n+1)A\cos(n+2)A = \cos A$

Ex: Prove That $\frac{\tan x + \tan y}{\tan x - \tan y} = \frac{\sin(x+y)}{\sin(x-y)}$

Ex: If $A + B = \frac{\pi}{4}$ then prove that $(1 + \tan A)(1 + \tan B) = 2$

Ex: Prove that $\frac{\cos 90^\circ + \sin 90^\circ}{\cos 90^\circ - \sin 90^\circ} = \tan 54^\circ$

Ex. Prove that $\sin^4 \frac{\pi}{8} + \sin^4 \frac{3\pi}{8} + \sin^4 \frac{5\pi}{8} + \sin^4 \frac{7\pi}{8} = \frac{3}{2}$

Ex. Prove that $\cot A - \operatorname{cosec} 2A = \cot 2A$

Ex. Prove that $\tan A(1 + \sec 2A) = \tan 2A$

Ex. Find the value of $2\sin 67\frac{1}{2}^\circ \cos 22\frac{1}{2}^\circ$

Ex. Prove that $\tan 75^\circ + \cot 75^\circ = 4$

Ex. Prove that $\tan 70^\circ = \tan 20^\circ + 2 \tan 50^\circ$

Ex: .Prove that $\frac{\sin 5x + \sin 3x}{\cos 5x + \cos 3x} = \tan 4x$

Ex: If $\tan A = x \tan B$ then prove that $\frac{\sin(A-B)}{\sin(A+B)} = \frac{x-1}{x+1}$

Ex: If $A + B = \frac{\pi}{4}$ prove that $(1 + \tan A)(1 + \tan B) = 2$

Ex: If $\tan A - \tan B = x$ and $\cot B - \cot A = y$ then prove that $\cot(A - B) = \frac{1}{x} + \frac{1}{y}$

Ex: Prove that $\frac{\cos 4x + \cos 3x + \cos 2x}{\sin 4x + \sin 3x + \sin 2x} = \cot 3x$

Ex: Prove that $\frac{1 + \sin x - \cos x}{1 + \sin x + \cos x} = \tan \frac{x}{2}$

Ex: Prove that $\frac{\cos x}{1 + \sin x} = \tan \left(\frac{\pi}{4} - \frac{x}{2}\right)$

Ex: Prove that $\cos^4 \frac{\pi}{8} + \cos^4 \frac{3\pi}{8} + \cos^4 \frac{5\pi}{8} + \cos^4 \frac{7\pi}{8} = \frac{3}{2}$

Ex: prove that $\frac{1 + \sin 2x + \cos 2x}{1 + \sin 2x - \cos 2x} = \cot x$

INVERSE TRIGONOMETRIC FUNCTION

Inverse Function : A function $f:X \rightarrow Y$ is said to be invertible (i.e f^{-1} exist) if it is bijective i.e one-one and on-to.

Here X is the domain and Y is the range of the function ,so $f^{-1}:Y \rightarrow X$.

If $y = f(x) \Rightarrow x = f^{-1}(y)$,Here the domain is Y and range is X.

Now consider a trigonometric function $y = \sin x$ which is not bijective in R.

So we have to restrict its domain in $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ corresponding to the range $[-1,1]$ then the function $\sin x$ is bijective .

Now $\sin: \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] \rightarrow [-1,1]$ is bijective and its inverse will exist,which can be written as $\sin^{-1}: [-1,1] \rightarrow \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$

Hence $y = \sin x \Rightarrow x = \sin^{-1}y$

Note : $\sin^{-1}x \neq (\sin x)^{-1} = \frac{1}{\sin x}$

Note: $f^{-1}Of(x) = f^{-1}f(x) = f^{-1}(y) = x$

$fOf^{-1}(y) = f(f^{-1}(y)) = f(x) = y$

Let us consider a function $y = \sin x$ then its inverse exist and which is written as $x = \sin^{-1}y$

Function	Domain	Range(Principal value)
\sin^{-1}	$[-1,1]$	$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$
\cos^{-1}	$[-1,1]$	$[0, \pi]$
\tan^{-1}	R	$\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$
\cot^{-1}	R	$(0, \pi)$
\cosec^{-1}	$R - (-1,1)$	$\left[-\frac{\pi}{2}, \frac{\pi}{2}\right] - \{0\}$
\sec^{-1}	$R - (-1,1)$	$[0, \pi] - \left\{\frac{\pi}{2}\right\}$

Formula-1

$$\sin(\sin^{-1}x) = x \quad x \in [-1,1]$$

$$\cos(\cos^{-1}x) = x \quad x \in [-1,1]$$

$$\tan(\tan^{-1}x) = x \quad x \in R$$

$$\cot(\cot^{-1}x) = x \quad x \in R$$

$$\sec(\sec^{-1}x) = x \quad x \in R - (-1,1)$$

$$\cosec(\cosec^{-1}x) = x, x \in R - (-1,1)$$

Formula-2

$$\sin^{-1}(\sin x) = x \quad x \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$$

$$\cos^{-1}(\cos x) = x \quad x \in [0, \pi]$$

$$\tan^{-1}(\tan x) = x \quad x \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$$

$$\cot^{-1}(\cot x) = x \quad x \in (0, \pi)$$

$$\sec^{-1}(\sec x) = x \quad x \in [0, \pi] - \left\{\frac{\pi}{2}\right\}$$

$$\cosec^{-1}(\cosec x) = x \quad x \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] - \{0\}$$

Formula-3

$$\sin^{-1}x = \cosec^{-1}\frac{1}{x} \quad , \cosec^{-1}x = \sin^{-1}\frac{1}{x} \quad , \cos^{-1}x = \sec^{-1}\frac{1}{x} \quad , \sec^{-1}x = \cos^{-1}\frac{1}{x}$$

$$\tan^{-1}x = \cot^{-1}\frac{1}{x} \quad , \cot^{-1}x = \tan^{-1}\frac{1}{x}$$

Formula-4

$$\sin^{-1}x + \cos^{-1}x = \frac{\pi}{2}, \quad x \in [-1, 1]$$

$$\tan^{-1}x + \cot^{-1}x = \frac{\pi}{2}, \quad x \in R$$

$$\sec^{-1}x + \cosec^{-1}x = \frac{\pi}{2}, \quad x \in R - (1, 1)$$

Formula-5

$$\sin^{-1}(-x) = -\sin^{-1}x \quad , \cos^{-1}(-x) = \pi - \cos^{-1}x \quad , \tan^{-1}(-x) = -\tan^{-1}x \quad ,$$

$$\cot^{-1}(-x) = -\cot^{-1}x \quad , \sec^{-1}(-x) = \pi - \sec^{-1}x \quad , \cosec^{-1}(-x) = -\cosec^{-1}x$$

Formula-6

$$\sin^{-1}x = \cos^{-1}\sqrt{1-x^2} = \tan^{-1}\frac{x}{\sqrt{1-x^2}} = \cot^{-1}\frac{\sqrt{1-x^2}}{x} = \sec^{-1}\frac{1}{\sqrt{1-x^2}} = \cosec^{-1}\frac{1}{x}$$

$$\cos^{-1}x = \sin^{-1}\sqrt{1-x^2} = \cot^{-1}\frac{x}{\sqrt{1-x^2}} = \tan^{-1}\frac{\sqrt{1-x^2}}{x} = \cosec^{-1}\frac{1}{\sqrt{1-x^2}} = \sec^{-1}\frac{1}{x}$$

$$\tan^{-1}x = \sec^{-1}\sqrt{1+x^2} = \sin^{-1}\frac{x}{\sqrt{1+x^2}} = \cosec^{-1}\frac{\sqrt{1+x^2}}{x} = \cos^{-1}\frac{1}{\sqrt{1+x^2}} = \cot^{-1}\frac{1}{x}$$

Formula-7

$$\tan^{-1}x + \tan^{-1}y = \tan^{-1}\frac{x+y}{1-xy}$$

$$\tan^{-1}x - \tan^{-1}y = \tan^{-1}\frac{x-y}{1+xy}$$

Formula-8

$$\sin^{-1}x + \sin^{-1}y = \sin^{-1}\left\{x\sqrt{1-y^2} + y\sqrt{1-x^2}\right\}$$

$$\sin^{-1}x - \sin^{-1}y = \sin^{-1}\left\{x\sqrt{1-y^2} - y\sqrt{1-x^2}\right\}$$

$$\cos^{-1}x + \cos^{-1}y = \sin^{-1}\left\{xy - \sqrt{1-x^2}\sqrt{1-y^2}\right\}$$

$$\cos^{-1}x - \cos^{-1}y = \sin^{-1}\left\{xy + \sqrt{1-x^2}\sqrt{1-y^2}\right\}$$

Formula-9

$$2\tan^{-1}x = \tan^{-1}\frac{2x}{1-x^2} = \sin^{-1}\frac{2x}{1+x^2} = \cos^{-1}\frac{1-x^2}{1+x^2}$$

Formula-10

$$2\sin^{-1}x = \sin^{-1}2x\sqrt{1-x^2}$$

$$2\cos^{-1}x = \cos^{-1}(2x^2 - 1) = \sin^{-1}(1 - 2x^2)$$

Formula-11

$$3\sin^{-1}x = \sin^{-1}\{3x - 4x^3\}$$

$$3\cos^{-1}x = \cos^{-1}\{4x^3 - 3x\}$$

$$3\tan^{-1}x = \tan^{-1}\left\{\frac{3x-x^3}{1-3x^2}\right\}$$

Assignment-1

1. Find the principal value of $\sin^{-1}\left(-\frac{1}{2}\right)$, $\cos^{-1}\left(-\frac{1}{2}\right)$, $\tan^{-1}(-\sqrt{3})$ and $\sec^{-1}\left(\frac{2}{\sqrt{3}}\right)$

2. Evaluate $\sin^{-1}\left(\sin\frac{2\pi}{3}\right)$, $\cos^{-1}\left(\cos(-\frac{\pi}{8})\right)$, $\sin^{-1}\left(\sin\frac{5\pi}{6}\right)$, $\cos^{-1}\left(\cos\frac{4\pi}{3}\right)$ and $\tan^{-1}\left(\frac{3\pi}{4}\right)$

3. True or False

$$(a) \cos\left(\cos^{-1}\frac{5}{13}\right) \quad (b) \tan(\tan^{-1}3) \quad (c) \sin\left(\sin^{-1}\frac{3}{2}\right)$$

4. True or False

$$(a) \sin^{-1}\frac{1}{2} + \cos^{-1}\frac{1}{2} = \frac{\pi}{2} \quad (b) \tan^{-1}5 + \cot^{-1}5 = \frac{\pi}{2} \quad (c) \sec^{-1}\frac{1}{4} + \operatorname{cosec}^{-1}\frac{1}{4} = \frac{\pi}{2}$$

5. Find the value of

$$(a) \sin\cos^{-1}\tan\sec^{-1}\sqrt{2} \quad (b) \cos\tan^{-1}\cot\cos^{-1}\frac{\sqrt{3}}{2} \quad (c) \cos\tan^{-1}\cot\sin^{-1}x \quad (d) \sin(\tan^{-1}x + \tan^{-1}\frac{1}{x})$$

$$6. \text{ Prove that } \tan^{-1}\frac{2}{11} + \tan^{-1}\frac{7}{24} = \tan^{-1}\frac{1}{2}$$

$$7. \text{ Prove that } \tan^{-1}\frac{1}{2} + \tan^{-1}\frac{1}{5} + \tan^{-1}\frac{1}{8} = \frac{\pi}{4}$$

$$8. \text{ Prove that } \tan^{-1}\frac{1}{7} + \tan^{-1}\frac{1}{5} + \tan^{-1}\frac{1}{8} + \tan^{-1}\frac{1}{3} = \frac{\pi}{4}$$

9. Prove that $\tan^{-1}\frac{3}{4} + \tan^{-1}\frac{3}{5} - \tan^{-1}\frac{8}{9} = \frac{\pi}{4}$

10. Find the value of $\tan\left(\frac{\pi}{4} + \tan^{-1}\frac{3}{4}\right)$

11. Prove that $\sin^{-1}\frac{5}{13} + \sin^{-1}\frac{3}{5} = \sin^{-1}\frac{56}{65}$

12. Prove that $\sin^{-1}\frac{4}{5} + \sin^{-1}\frac{5}{13} + \sin^{-1}\frac{16}{65} = \frac{\pi}{2}$

13. Prove that $\cos^{-1}\frac{4}{5} - \cos^{-1}\frac{15}{17} = \cos^{-1}\frac{84}{85}$

14. Prove that $\cos^{-1}\frac{3}{5} + \cos^{-1}\frac{12}{13} + \cos^{-1}\frac{63}{65} = \frac{\pi}{2}$

15. Find the value of $\cos\left(\sin^{-1}\frac{5}{13} + \sin^{-1}\frac{3}{5}\right)$

Assignment-2:

1. Prove that $4\tan^{-1}\frac{1}{5} - \tan^{-1}\frac{1}{70} + \tan^{-1}\frac{1}{99} = \frac{\pi}{4}$

2. Prove that $\sec^{-1}\frac{5\sqrt{2}}{7} + 2\tan^{-1}\frac{1}{5} + 2\tan^{-1}\frac{1}{8} = \frac{\pi}{4}$

3. Evaluate $\sin(2\sin^{-1}0.6)$, $\sin(2\sin^{-1}0.8)$, and $\sin(3\sin^{-1}0.4)$

4. Prove that $2\tan^{-1}\frac{1}{2} + \tan^{-1}\frac{1}{7} = \tan^{-1}\frac{31}{17}$

5. Prove that $\tan\left\{\frac{\pi}{4} + \frac{1}{2}\cos^{-1}\frac{a}{b}\right\} + \tan\left\{\frac{\pi}{4} - \frac{1}{2}\cos^{-1}\frac{a}{b}\right\} = \frac{2b}{a}$

6. Prove that $\sec^2(\tan^{-1}2) + \operatorname{cosec}^2(\cot^{-1}3) = 15$

7. Evaluate $\tan\left(2\tan^{-1}\frac{1}{5}\right)$

8. Evaluate $\tan\left(2\tan^{-1}\frac{1}{5} - \frac{\pi}{4}\right)$

9. Evaluate $\tan\left(\frac{1}{2}\cos^{-1}\frac{\sqrt{5}}{3}\right)$

10. Solve $\tan^{-1}2x + \tan^{-1}3x = \frac{\pi}{4}$

11. Solve $\sin^{-1}(1-x) - 2\sin^{-1}x = \frac{\pi}{2}$

12. Solve $\sin^{-1}x + \sin^{-1}2x = \frac{\pi}{3}$

13. Solve $\cos^{-1}x + \sin^{-1}\frac{x}{2} = \frac{\pi}{6}$

14. Prove that $\sin^2(\sin^{-1}x + \sin^{-1}y + \sin^{-1}z) = \cos^2(\cos^{-1}x + \cos^{-1}y + \cos^{-1}z)$

15. If $\sin^{-1}x + \sin^{-1}y + \sin^{-1}z = \pi$ then prove that $x\sqrt{1-x^2} + y\sqrt{1-y^2} + z\sqrt{1-z^2} = 2xyz$